



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

# THE AMERICAN MATHEMATICAL MONTHLY.

Entered at the Post-office at Springfield, Missouri, as Second-class Mail Matter.

VOL. V.

MARCH, 1898.

No. 3.

## NON-EUCLIDEAN GEOMETRY: HISTORICAL AND EXPOSITORY.

By GEORGE BRUCE HALSTED, A. M. (Princeton); Ph. D. (Johns Hopkins); Member of the London Mathematical Society; and Professor of Mathematics in the University of Texas, Austin, Texas.

[Continued from January Number.]

**PROPOSITION XXXII.** *Now I say there is (in hypothesis of acute angle) a certain determinate acute angle  $BAX$  drawn under which  $AX$  (Fig. 33.) only at an infinite distance meets  $BX$ , and thus is a limit in part from within, in part from without; on the one hand of all those which under lesser acute angles meet the aforesaid  $BX$  at a finite distance; on the other hand also of the others which under greater acute angles, even to a right angle inclusive, have a common perpendicular in two distinct points with  $BX$ .*

**PROOF.** First it holds (from Cor. II. after Proposition XXIX.) that no determinate acute angle will be the greatest of all drawn under which a straight from the point  $A$  meets the aforesaid  $BX$  at a finite distance

Secondly, it holds in like manner that (in hypothesis of acute angle) no acute angle will be the least of all drawn under which a straight has a common perpendicular in two distinct points with  $BX$ ; since indeed (from what precedes) there can be no determinate limit, such that there cannot be found, under a lesser angle




Fig. 33.

be no determinate limit, such that there cannot be found, under a lesser angle

constituted at the point  $A$ , a common perpendicular in two distinct points, which is less than any assignable length  $R$ .

And hence follows thirdly, that (in this hypothesis) there must be a certain determinate acute angle  $BAX$ , drawn under which  $AX$  so approaches ever more to  $BX$ , that only at an infinite distance does it meet it.

But further that this  $AX$  is a limit in part from within in part from without of each of the aforesaid classes of straights is proved thus. First, it agrees with those straights which meet  $BX$  at a finite distance since it also finally meets; but it differs, because it meets only at an infinite distance.

But secondly it also agrees, and at the same time differs from those straights which have a common perpendicular in two distinct points with  $BX$ ; because it also has a common perpendicular with  $BX$ ; but in one and the same point  $X$  infinitely distant. But this latter ought to be considered demonstrated in Proposition XXVIII., as I point out in its corollary.

Therefore it holds, that (in the hypothesis of acute angle) there will be a certain determinate acute angle  $BAX$ , drawn under which  $AX$  only at an infinite distance meets  $BX$ , and thus is a limit in part from within, in part from without; on the one hand of all those which under lesser acute angles meet the aforesaid  $BX$  at a finite distance; on the other hand also of the others which under greater acute angles, even to a right angle inclusive, have a common perpendicular in two distinct points with  $BX$ . Quod erat etc.

[To be Continued.]

---

## ON THE BEST METHOD OF SOLVING THE MARKINGS OF JUDGES OF CONTESTS.

By F. R. MOULTON.

---

The fact that many different methods are in use for deciding from the markings of judges the relative standing of the participants in oratorical and similar contests, and that several different methods have been stated to be the best by Professors of Mathematics in some of our colleges, may be taken as the excuse for this paper. It is questionable whether such a problem can be solved by perfectly rigorous mathematical processes, but it has seemed that a method similar to that of Hagen in the theory of probability may be applied with advantage. Let us agree to adopt the following hypotheses.

- (1) The judges mark each contestant independently and by the same scale.
- (2) There is a true marking for each contestant as compared to a fixed ideal.
- (3) The deviation of each judge's markings from these true markings, are the result of a very great many influences, such as, the inherited inclinations re-